

PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Performance of Low-Risk Criteria in the Evaluation of Young Infants With Fever: Review of the Literature

Anna R. Huppler, Jens C. Eickhoff and Ellen R. Wald

Pediatrics 2010;125:228-233; originally published online Jan 18, 2010;

DOI: 10.1542/peds.2009-1070

The online version of this article, along with updated information and services, is
located on the World Wide Web at:

<http://www.pediatrics.org/cgi/content/full/125/2/228>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2010 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



Performance of Low-Risk Criteria in the Evaluation of Young Infants With Fever: Review of the Literature



WHAT'S KNOWN ON THIS SUBJECT: Fever in neonates is common. The rate of SBIs in young infants may be as high as 12%. Low-risk criteria have been developed to aid in management decisions for well-appearing, febrile young infants.



WHAT THIS STUDY ADDS: Although the total risk of SBI in febrile young infants in this review was 10.9%, low-risk criteria allowed 30% of these patients to be treated safely without empiric antibiotic therapy.

abstract

FREE

OBJECTIVE: The goal was to determine the performance of low-risk criteria for serious bacterial illnesses (SBIs) in febrile infants in prospective studies in which empiric antibiotic treatment was withheld, compared with studies (prospective and retrospective) in which empiric antibiotic treatment was administered.

METHODS: A search of the English-language literature was undertaken by using a PubMed database and reference lists of relevant studies of fever, low-risk criteria, and SBIs. Studies of infants >90 days of age, infants with specific infections, or infants with additional risk factors for infection were excluded. Publications were categorized as retrospective, prospective with empiric antibiotic treatment for all patients, or prospective with antibiotics withheld. The relative risk of SBI in high-risk versus low-risk patients was determined for pooled data in each category. The rates of SBIs in low-risk patients in each category were compared.

RESULTS: Twenty-one studies met the inclusion criteria. In prospective studies in which patients were cared for without empiric antibiotic treatment, 6 patients assigned to the low-risk category had SBIs; all recovered uneventfully. The rate of SBIs in these low-risk patients was 0.67%. The relative risk of SBIs in high-risk versus low-risk patients in these studies was 30.56 (95% confidence interval: 7.0–68.13). The rate of SBIs in low-risk patients in all studies was 2.23%. The rate of SBIs in low-risk patients in the prospective studies without empiric antibiotic treatment was significantly different from the rate in all other studies (0.67% vs 2.71%; $P = .01$).

CONCLUSIONS: Low-risk criteria perform well in prospective studies in which empiric antibiotic treatment is withheld. These criteria allow ~30% of young febrile infants to be observed without antibiotic treatment, thus avoiding unnecessary hospitalization, nosocomial infection, injudicious use of antibiotics, and adverse effects of antibiotics. *Pediatrics* 2010;125:228–233

AUTHORS: Anna R. Huppler, MD,^a Jens C. Eickhoff, PhD,^{b,c} and Ellen R. Wald, MD^a

Departments of ^aPediatrics and ^bBiostatistics and Medical Informatics, University of Wisconsin-Madison, Madison, Wisconsin; and ^cDepartment of Statistics and Clinical Sciences, Colorado State University, Fort Collins, Colorado

KEY WORDS

fever, infant, bacteremia, meningitis, urinary tract infection

ABBREVIATIONS

SBI—serious bacterial illness

CI—confidence interval

RR—relative risk

UTI—urinary tract infection

www.pediatrics.org/cgi/doi/10.1542/peds.2009-1070

doi:10.1542/peds.2009-1070

Accepted for publication Aug 28, 2009

Address correspondence to Anna R. Huppler, MD, University of Wisconsin-Madison, Department of Pediatrics, 600 Highland Ave, Madison, WI 53792. E-mail: hupplerar@upmc.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2010 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: *The authors have indicated they have no financial relationships relevant to this article to disclose.*

Fever in very young infants is a common and important problem. Neonates have unique vulnerabilities to infection because of their immature immune systems and incomplete barriers to invasion. The rate of serious bacterial illnesses (SBIs) in young febrile infants has been reported to be between 8.5% and 12%.^{1–3}

Before 1985, it was recommended that all febrile infants (<60 days of age) be hospitalized and treated with parenteral antibiotic therapy after a full sepsis evaluation, because various criteria used to identify “high-risk” infants were insufficiently sensitive to identify all infants with SBIs.^{4–6} However, the approach of hospitalizing and treating all febrile young infants with empiric antibiotic therapy had the disadvantages of unnecessary hospitalizations, nosocomial infections, injudicious use of antibiotics, emergence of resistant bacteria, and adverse effects of antibiotics.^{7,8}

During the late 1980s and early 1990s, investigators changed strategies and attempted to identify febrile infants who were at low risk for SBIs and might be treated with close observation (inpatient or outpatient) without antibiotic treatment.^{3,9–16} After the development of various low-risk criteria, several groups undertook studies to validate the criteria in their own populations.^{17–27} The designs of those studies alternated between retrospective and prospective, with variable approaches to empiric use of ceftriaxone. We hypothesized that prospective studies that implemented a strategy of observation without antibiotic treatment for low-risk infants, thereby relying on meticulous evaluation and careful decision-making, would show significantly better performance of low-risk criteria than would studies that continued to treat all patients (prospectively or retrospectively), regardless of risk stratification.

METHODS

We searched the English-language literature for original articles studying low-risk criteria for SBIs in febrile infants between 0 and 90 days of age. A National Library of Medicine PubMed database search for articles was performed by using combinations of the following search terms: “low risk criteria,” “criteria,” or “risk”; “serious bacterial illness,” “serious bacterial infection,” “bacterial infection,” or “bacteremia”; and “fever” or “fever without source”. Studies were limited to humans, infants, and publication after 1985.

The search criteria identified 740 articles. Bibliographic references from the relevant studies (original research and review articles) were reviewed for additional citations. Articles were excluded on the basis of abstracts if they focused on infants >90 days of age, patients with underlying conditions (such as patients with sickle cell disease, neutropenia, malignancy, central vascular catheters, or other immunocompromised states), or patients with a single focus of bacterial infection (eg, urinary tract infection [UTI] or meningitis). Sixty-three publications were reviewed carefully for inclusion criteria, definition of SBIs, and evaluations performed for study patients. The components of the low-risk criteria used in each study were reviewed and compared with the original Rochester Criteria⁹ (Tables 1 and 2). Studies with overlapping subjects were excluded.

Articles were grouped according to type of study, and total patients were aggregated. The categories of studies were prospective without antibiotic treatment of low-risk infants, prospective with antibiotic treatment of low-risk infants, and retrospective.

Portions of studies using different management plans for low-risk infants were classified accordingly. Data were

TABLE 1 Rochester Criteria for Infants at Low Risk of SBIs^{9,27}

| |
|---|
| 1. Previously healthy term infant without perinatal complications and with no previous antibiotic treatment |
| 2. Normal physical examination findings (including no otitis media) |
| 3. White blood cell count: 5000–15 000 cells per mm ³ |
| 4. Band count: <1500 cells per mm ³ |
| 5. Urinalysis: <10 white blood cells per high-power field in centrifuged catheterized specimen |
| No comment about stool |

Reference 9 established the original Rochester criteria. Reference 27 used identical laboratory criteria and slightly different inclusion criteria.

extracted from the tables and text of each study and included the number of low-risk infants, the number of high-risk infants, the number of SBIs in each group, types of SBIs, and the outcomes of low-risk infants with SBIs. The random-effects method described by DerSimonian and Laird²⁸ was used to estimate overall relative risks (RRs) and corresponding 95% confidence intervals (CIs). This method accounts for the heterogeneity of studies through a statistical parameter representing interstudy variation.

Heterogeneity between studies was assessed by using the *Q* statistic, as well as graphic techniques. The β -binomial model for overdispersed data²⁹ was used to estimate pooled rates of SBIs for low- and high-risk patients. Comparison of pooled rates between groups was performed by using a likelihood ratio test. Linear regression analysis was used to evaluate the trend in the proportions of infants designated as being at low risk over time.

RESULTS

Twenty-one studies met inclusion criteria; 14 were prospective studies (Tables 3 and 4) and 7 were retrospective (Table 5).^{3,9–27,30} Among the prospective studies, 9 treated low-risk patients empirically with antibiotics, 4 monitored low-risk patients without antibiotic treatment, and 1 used both strategies.

TABLE 2 Variations of Rochester Criteria

| Type of Low-Risk Criteria | Differences From Original Rochester Criteria | References |
|---------------------------|---|--------------------|
| Rochester 2 | If diarrhea, ≤ 5 –10 WBCs per high-power field in stool | 10, 17, 18, 20, 30 |
| Modified Rochester | Normal inflammatory markers (C-reactive protein levels or ESR) | 13, 19, 21 |
| Milwaukee | CSF: < 10 WBCs per mm^3 ; WBC count: $< 15\,000$ cells per mm^3 (no band criteria); urinalysis: ≤ 5 –10 WBCs per high-power field, no bacteria; urine dipstick: negative LE/nitrite | 14 |
| Philadelphia | Infant observation score ³⁵ : < 10 ; WBC count: $< 15\,000$ cells per mm^3 (no band criteria); urinalysis: < 10 WBCs per high-power field, few or no bacteria; CSF: < 8 WBCs per mm^3 , no bacteria, nonbloody | 11 |
| Philadelphia 2 | WBC count: $< 15\,000$ cells per mm^3 ; band/neutrophil ratio: < 0.2 ; CSF: < 8 WBCs per mm^3 , no bacteria, nonbloody | 20, 22–25 |
| Boston | WBC count: $< 20\,000$ cells per mm^3 (no band criteria); CSF: < 10 WBCs per mm^3 ; urinalysis: < 10 WBCs per high-power field, no LE | 12, 24 |
| Pittsburgh | Enhanced urinalysis: ≤ 9 WBCs per mm^3 , negative Gram stain results; CSF: ≤ 5 WBCs per mm^3 , negative Gram stain results (if < 6 wk) | 26 |
| Impression of sepsis | “Not ill” or negative clinical impression of sepsis (history, physical examination, ESR of < 30 mm/h, WBC count of $< 15\,000$ cells per mm^3) | 3, 15, 16 |

WBC indicates white blood cell; ESR, erythrocyte sedimentation rate; CSF, cerebrospinal fluid; LE, leukocyte esterase. The infant observation score includes 5 observations, that is, quality of cry, reaction to parent stimulation and state variation, color, hydration, and response to social overtures, each scored on a scale of 1 to 5.³¹

The type of low-risk criteria was categorized, with 10 studies using variations of the Rochester criteria, 6 studies using Philadelphia criteria (2 comparing Rochester criteria with Philadelphia criteria), 2 studies using Boston criteria, 1 study using Pittsburgh criteria, 1 study using Milwaukee criteria, and 3 studies using “clinical impression of sepsis” or “not ill” descriptions (Tables 1 and 2). Studies were performed between 1979 and 1999. One study¹² included only children in the low-risk category, without a high-risk comparison group. Fever

was defined as a temperature of $> 38.2^\circ\text{C}$ in 1 study,¹¹ $> 38.1^\circ\text{C}$ in 3 studies,^{15,20,27} and $> 38.0^\circ\text{C}$ in the rest of the studies.

SBLs included bacteremia, meningitis, bacterial diarrhea, pneumonia, and UTIs.³¹ The defining characteristics for each type of infection varied slightly between the studies, most notably in the definition of UTI. Two studies accepted bagged urine specimens for culture, resulting in 8 diagnoses of UTIs in low-risk patients.^{15,19} Ten studies diagnosed UTIs in patients with

catheterized urine cultures with $< 50\,000$ colony-forming units of a single organism per milliliter.^{11,12,14,17,18,20–23,25} Four studies provided no microbiologic definition of UTI or description of urine collection methods.^{3,15,16,30} Another variation was the inclusion of pneumonia as a SBL without microbiologic documentation. All infants studied were between 0 and 90 days of age. Six studies included only infants in the first month of life, and 4 studies excluded infants in the first month of life. The majority of studies were limited to the first 2 months of life, with 6 studies extending to 3 months.

A summary of the prospective studies in which patients underwent observation alone is shown in Table 3.^{10,11,18,19,22} A total of 1858 patients were included, and 870 were classified as being at low risk. Six patients with SBLs were missed, including 2 patients with bacteremia and 4 patients with UTIs. The first child with bacteremia received parenteral antibiotic treatment within 24 hours after presentation and recovered uneventfully.¹¹ The identity of the organism was not noted. The second child had a blood culture positive for *Yersinia enterocolitica* and fared well after treatment.¹⁸ The authors noted that a confirmatory culture was not performed before initiation of treatment. Three urine specimens obtained with sterile technique yielded either group B streptococcus or *Escherichia coli*.¹⁸ Those patients fared well with treatment. One urine specimen collected by bag yielded *E coli*.¹⁹

TABLE 3 Prospective Studies of Performance of Low-Risk Criteria for SBLs in Young Infants in Which Antibiotics Were Withheld From Low-Risk Patients

| Year and Reference | Criteria Type | Age, d | No. of Patients | No. of High-Risk Patients | Cases of SBL in High-Risk Patients | Rate of SBLs in High-Risk Patients, % | No. of Low-Risk Patients | Cases of SBL in Low-Risk Patients | Rate of SBLs in Low-Risk Patients, % |
|--------------------|--------------------|--------|-----------------|---------------------------|------------------------------------|---------------------------------------|--------------------------|-----------------------------------|--------------------------------------|
| 1988 ¹⁰ | Rochester 2 | 0–56 | 236 | 88 | 21 | 23.9 | 148 | 0 | 0.00 |
| 1993 ¹¹ | Philadelphia | 29–56 | 747 | 460 | 64 | 13.9 | 287 | 1 | 0.35 |
| 1994 ¹⁸ | Rochester 2 | 0–60 | 203 | | | | 203 | 4 | 1.97 |
| 1997 ¹⁹ | Modified Rochester | 4–28 | 250 | 119 | 40 | 33.6 | 131 | 1 | 0.76 |
| 1999 ²² | Philadelphia 2 | 29–60 | 422 | 321 | 43 | 13.4 | 101 | 0 | 0.00 |
| Total | | | 1858 | 988 | 168 | 20.6 | 870 | 6 | 0.67 |

TABLE 4 Prospective Studies of Performance of Low-Risk Criteria for SBIs in Young Infants With Empiric Use of Antibiotics for All Patients

| Year and Reference | Criteria Type | Age, d | No. of Patients | No. of High-Risk Patients | Cases of SBI in High-Risk Patients | Rate of SBIs in High-Risk Patients, % | No. of Low-Risk Patients | Cases of SBI in Low-Risk Patients | Rate of SBIs in Low-Risk Patients, % |
|--------------------|--------------------------------|---------|-----------------|---------------------------|------------------------------------|---------------------------------------|--------------------------|-----------------------------------|--------------------------------------|
| 1985 ⁹ | Rochester | 0–90 | 233 | 89 | 22 | 24.7 | 144 | 1 | 0.69 |
| 1987 ¹⁶ | Impression of sepsis | 0–56 | 97 | 36 | 5 | 13.9 | 61 | 2 | 3.28 |
| 1988 ¹⁵ | Impression of sepsis | 0–14 | 35 | 11 | 4 | 36.4 | 24 | 2 | 8.33 |
| 1992 ¹² | Boston | 28–89 | 503 | | | | 503 | 27 | 5.37 |
| 1993 ¹⁴ | Milwaukee | 28–56 | 534 | 391 | 23 | 5.9 | 143 | 1 | 0.70 |
| 1994 ¹⁸ | Rochester 2 | 0–60 | 802 | 494 | 61 | 12.3 | 308 | 1 | 0.32 |
| 1994 ¹³ | Modified Rochester | 0–31 | 254 | 120 | 37 | 30.8 | 134 | 8 | 5.97 |
| 2004 ³⁰ | Rochester 2 | 0–90 | 1378 | 922 | 118 | 12.8 | 456 | 12 | 2.63 |
| 2005 ²⁰ | Rochester 2 and Philadelphia 2 | 0–56 | 259 | 186 | 63 | 33.9 | 73 | 2 | 2.74 |
| 2007 ²¹ | Modified Rochester | Neonate | 386 | 220 | 107 | 48.6 | 166 | 1 | 0.60 |
| Total | | | 4481 | 2469 | 440 | 23.8 | 2012 | 57 | 2.71 |

TABLE 5 Retrospective Studies of Performance of Low-Risk Criteria for SBIs in Young Infants

| Year and Reference | Criteria Type | Age, d | No. of Patients | No. of High-Risk Patients | Cases of SBI in High-Risk Patients | Rate of SBIs in High-Risk Patients, % | No. of Low-Risk Patients | Cases of SBI in Low-Risk Patients | Rate of SBIs in Low-Risk Patients, % |
|--------------------|---------------------------|--------|-----------------|---------------------------|------------------------------------|---------------------------------------|--------------------------|-----------------------------------|--------------------------------------|
| 1986 ²⁷ | Rochester | 0–90 | 117 | 47 | 4 | 8.5 | 70 | 3 | 4.29 |
| 1990 ³ | Impression of sepsis | 0–91 | 443 | 222 | 48 | 21.6 | 221 | 5 | 2.26 |
| 1997 ¹⁷ | Rochester 2 | 0–29 | 119 | 71 | 19 | 26.8 | 48 | 3 | 6.25 |
| 1997 ²⁵ | Philadelphia 2 | 3–90 | 492 | 196 | 52 | 26.5 | 296 | 8 | 2.70 |
| 1999 ²³ | Philadelphia 2 | 3–28 | 254 | 145 | 27 | 18.6 | 109 | 5 | 4.59 |
| 2000 ²⁴ | Boston and Philadelphia 2 | 1–28 | 372 | 141 | 37 | 26.2 | 231 | 8 | 3.46 |
| 2001 ²⁶ | Pittsburgh | 0–60 | 404 | 277 | 41 | 14.8 | 127 | 0 | 0.00 |
| Total | | | 2201 | 1099 | 228 | 19.8 | 1102 | 32 | 2.70 |

TABLE 6 Comparison of Performance of Low-Risk Criteria for SBIs in Young Infants, According to Study Design

| Type of Study | Total No. of Patients | No. of Patients With SBIs | No. of High-Risk Patients | Pooled Rate of SBIs in High-Risk Patients, Estimate (95% CI), % ^a | No. of Low-Risk Patients | Pooled Rate of SBIs in Low-Risk Patients, Estimate (95% CI), % ^a | RR of SBI in High-Risk vs Low-Risk Patients (95% CI) |
|---|-----------------------|---------------------------|---------------------------|--|--------------------------|---|--|
| Prospective, no antibiotic treatment | 1858 | 174 | 988 | 20.6 (9.4–31.8) | 870 | 0.67 (–0.04–1.30) | 30.56 (7.0–68.13) |
| Prospective, empiric antibiotic treatment | 4481 | 497 | 2469 | 23.8 (13.4–34.1) | 2012 | 2.71 (0.93–4.50) | 8.75 (2.29–15.21) |
| Retrospective | 2201 | 260 | 1099 | 19.8 (14.5–25.1) | 1102 | 2.70 (0.40–5.02) | 6.93 (3.10–10.75) |
| Retrospective and prospective, empiric antibiotic treatment | 6682 | 757 | 3568 | 22.3 (15.8–28.3) | 3114 | 2.71 (1.4–4.0) | 7.74 (3.82–11.67) |

^a Estimated by using the β -binomial model for overdispersed data.²⁸

In the rest of the studies summarized in Tables 4 and 5 (prospective studies in which patients received empiric antibiotic treatment and retrospective studies), 89 low-risk infants (2.71%) were diagnosed as having SBIs. This total included 2 cases of meningitis (1 with UTI and 1 with bacteremia), 22 cases of bacteremia (1 with gastroenteritis and 1 with osteomyelitis), 39 cases of UTI, and 14 cases of gastroenteritis. Twelve cases of SBI did not have a source identified.

Table 6 shows a comparison of the performance of the low-risk criteria for SBIs in young infants according to study design. The overall validity of the low-risk criteria in pooled studies was assessed by calculating the RR for SBI in high-risk versus low-risk infants. The RR reached statistical significance in all 3 categories of studies. In prospective studies with no empiric antibiotic treatment, the RR was 30.56 (95% CI: 7.0–68.13). In prospective studies with empiric antibiotic treat-

ment, the RR was 8.75 (95% CI: 2.29–15.21). In retrospective studies, the RR was 6.93 (95% CI: 3.10–10.75). Of importance, there was a statistically significant difference in the rates of SBIs in low-risk patients in the prospective studies with no empiric antibiotic treatment, compared with all other studies (P for difference = .010). When prospective studies with no empiric antibiotic treatment were compared with all other studies, no significant difference in the rate of SBIs in high-




FIGURE 1 Proportions of infants assigned to the low-risk category according to year of study initiation, with a statistically significant trend toward assignment of fewer infants to the low-risk category over time ($P = .001$).

risk patients was found (RR: 1.10; $P = .75$).

An analysis examining the proportion of infants designated as low risk in all studies showed a significant trend toward assigning fewer infants to the low-risk category over time ($\rho = -0.67$; $P = .001$). The slope parameter of the linear regression model of the proportions of infants designated as low risk by time (year) was -0.015 ($P < .001$). Early studies placed $\sim 60\%$ of children in the low-risk category. By the middle 1990s, only 30% of children were considered at low risk, as illustrated in Fig 1.

DISCUSSION

The impetus for this study was the apparent discrepancy in the rate of SBIs among low-risk febrile infants reported in the literature. Previous studies reported a range of rates of SBIs in low-risk infants of 0% to

8.3%.^{10,15,22,26} This can be explained, in part, by the different sets of criteria used by various investigators (Table 2).^{3,9–26,30} We showed that the rates of SBIs in low-risk patients in both retrospective studies and prospective studies using empiric antibiotic treatment were the same (2.7%) and were significantly different from the rate of SBIs in low-risk patients in prospective studies in which antibiotics were withheld (0.67%). We hypothesized that the low-risk criteria would function best when used in prospective studies in which low-risk patients underwent observation alone. Without reliance on empiric antibiotic treatment, it would be essential to capture all infants at risk of early deterioration in the high-risk group. Careful sample collection, as well as meticulous physical examination, excluded infants with SBIs from the low-risk group.

For physicians to rely on clinical algorithms, they must be convinced that patients will not be placed in jeopardy. The clinical predictability provided by low-risk criteria will be deemed satisfactory if the children identified as being at low risk either do not have a SBI or remain in stable condition until the SBI is recognized. In prospective trials that treated low-risk infants with observation alone, the low-risk criteria “missed” SBIs in 6 patients (bacteremia in 2 patients and UTIs in 4 patients). One of the cases involved a positive urine culture from a bagged specimen, which may represent a contaminant. These patients were treated with appropriate antibiotics when the cultures yielded positive results, and all recovered uneventfully. Therefore, careful application of these low-risk criteria was very effective in identifying children from whom empiric antibiotic therapy could be withheld.

Overall, data from 8540 infants were analyzed in this study. The total number of SBIs was 931, that is, 10.9%, consistent with rates commonly reported in the literature. With the use of low-risk criteria, $\sim 30\%$ of febrile infants can be identified as being at low risk for SBIs and can be treated with observation alone. These patients can be spared the negative effects that may be associated with empiric use of antibiotics, including cost, adverse effects of medications, development of resistant organisms, and psychosocial stresses on family dynamics. The criteria must be applied carefully to avoid misassignment of infants. Special attention should be given to the physical examination and medical history.

This study represents a comprehensive review of the literature that has evaluated the effectiveness of low-risk criteria in the identification of infants unlikely to have SBIs over the previous

23 years. The major weakness of this analysis is that the studies demonstrated minor variations in the low-risk criteria used and the age groups included. Although these small differ-

ences in demographic characteristics prohibit a “clean” comparison of all of the studies, we conclude that the observed difference in the performance of low-risk criteria was a function of

the study design. When low-risk criteria are applied prospectively after careful evaluation, clinicians should be confident in withholding empiric antibiotic treatment.

REFERENCES

- Bonadio WA. Evaluation and management of serious bacterial infections in the febrile young infant. *Pediatr Infect Dis J*. 1990; 9(12):905–912
- Levine DA, Platt SL, Dayan PS, et al. Risk of serious bacterial infection in young febrile infants with respiratory syncytial virus infections. *Pediatrics*. 2004;113(6): 1728–1734
- Wasserman GM, White CB. Evaluation of the necessity for hospitalization of the febrile infant less than three months of age. *Pediatr Infect Dis J*. 1990;9(3):163–169
- Caspe WB, Chamudes O, Louie B. The evaluation and treatment of the febrile infant. *Pediatr Infect Dis*. 1983;2(2):131–135
- Long SS. Approach to the febrile patient with no obvious focus of infection. *Pediatr Rev*. 1984;5(10):305–315
- Roberts KB, Brozy MS. Fever in the first 8 weeks of life. *Johns Hopkins Med J*. 1977; 141(1):9–13
- DeAngelis C, Joffe A, Wilson M, et al. Iatrogenic risks and financial costs of hospitalizing febrile infants. *Am J Dis Child*. 1983; 137(12):1146–1149
- Lieu TA, Baskin MN, Schwartz JS, Fleisher GR. Clinical and cost-effectiveness of outpatient strategies for management of febrile infants. *Pediatrics*. 1992;89(6):1135–1144
- Dagan R, Poell KR, Hall CB, Menegus MA. Identification of infants unlikely to have serious bacterial infection although hospitalized for suspected sepsis. *J Pediatr*. 1985; 107(6):855–860
- Dagan R, Sofer S, Phillip M, Shachak E. Ambulatory care of febrile infants younger than 2 months of age classified as being at low risk for having serious bacterial infections. *J Pediatr*. 1988;112(3):355–360
- Baker MD, Bell LM, Avner JR. Outpatient management without antibiotics in fever in selected infants. *N Engl J Med*. 1993;329(20): 1437–1441
- Baskin MN, O'Rourke EJ, Fleisher GR. Outpatient treatment of febrile infants 28 to 89 days of age with intramuscular administration of ceftriaxone. *J Pediatr*. 1992;120(1): 22–27
- Chiu C-H, Lin T-Y, Bullard MJ. Application of criteria identifying febrile outpatient neonates at low risk for bacterial infections. *Pediatr Infect Dis J*. 1994;13(11):946–949
- Bonadio WA, Hagen E, Rucka J, Shallow K, Stommel P, Smith D. Efficacy of a protocol to distinguish risk of serious bacterial infection in the outpatient evaluation of febrile young infants. *Clin Pediatr (Phila)*. 1993; 32(7):401–404
- Crain EF, Gershel JC. Which febrile infants younger than two weeks of age are likely to have sepsis? A pilot study. *Pediatr Infect Dis J*. 1988;7(8):561–564
- King JC, Berman ED, Wright PF. Evaluation of fever in infants less than 8 weeks old. *South Med J*. 1987;80(8):948–952
- Ferrera PC, Bartfield JM, Snyder HS. Neonatal fever: utility of the Rochester criteria in determining low risk for serious bacterial infections. *Am J Emerg Med*. 1997;15(3): 299–302
- Jaskiewicz JA, McCarthy CA, Richardson AC, et al. Febrile infants at low risk for serious bacterial infection: an appraisal of the Rochester criteria and implications for management. *Pediatrics*. 1994;94(3): 390–396
- Chiu C-H, Lin T-Y, Bullard MJ. Identification of febrile neonates unlikely to have bacterial infections. *Pediatr Infect Dis J*. 1997;16(1): 59–63
- Garra G, Cunningham SJ, Crain EF. Reappraisal of criteria used to predict serious bacterial illness in febrile infants less than 8 weeks of age. *Acad Emerg Med*. 2005; 12(10):921–925
- Marom R, Sakran W, Antonelli J, et al. Quick identification of febrile neonates with low risk for serious bacterial infection: an observational study. *Arch Dis Child Fetal Neonatal Ed*. 2007;92(1):F15–F18
- Baker MD, Bell LM, Avner JR. The efficacy of routine outpatient management without antibiotics of fever in selected infants. *Pediatrics*. 1999;103(3):627–631
- Baker MD, Bell LM. Unpredictability of serious bacterial illness in febrile infants from birth to 1 month of age. *Arch Pediatr Adolesc Med*. 1999;153(5):508–511
- Kadish HA, Loveridge B, Tobey J, Bolte RG, Cornell HM. Applying outpatient protocols in febrile infants 1–28 days of age: can the threshold be lowered? *Clin Pediatr (Phila)*. 2000;39(2):81–88
- Brik R, Hamissah R, Shehada N, Berant M. Evaluation of febrile infants under 3 months of age: is routine lumbar puncture warranted? *Isr J Med Sci*. 1997;33(2):93–97
- Herr SM, Wald ER, Pitetti RD, Choi SS. Enhanced urinalysis improves identification of febrile infants ages 60 days and younger at low risk for serious bacterial illness. *Pediatrics*. 2001;108(4):866–871
- Anbar RD, Richardson-de Corral V, O'Malley PJ. Difficulties in universal application of criteria identifying infants at low risk for serious bacterial infection. *J Pediatr*. 1986; 109(3):483–485
- DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials*. 1986;7(3): 177–188
- Young-Xu Y, Chan KA. Pooling overdispersed binomial data to estimate event rate. *BMC Med Res Methodol*. 2008;8(1):58
- Byington CL, Enriquez FR, Hoff C, et al. Serious bacterial infections in febrile infants 1 to 90 days old with and without viral infections. *Pediatrics*. 2004;113(6):1662–1666
- McCarthy PL, Sharpe MR, Spiesel SZ, et al. Observation scales to identify serious illness in febrile children. *Pediatrics*. 1982; 70(5):802–809

Performance of Low-Risk Criteria in the Evaluation of Young Infants With Fever: Review of the Literature

Anna R. Huppler, Jens C. Eickhoff and Ellen R. Wald

Pediatrics 2010;125:228-233; originally published online Jan 18, 2010;

DOI: 10.1542/peds.2009-1070

Updated Information & Services

including high-resolution figures, can be found at:
<http://www.pediatrics.org/cgi/content/full/125/2/228>

References

This article cites 31 articles, 14 of which you can access for free at:
<http://www.pediatrics.org/cgi/content/full/125/2/228#BIBL>

Post-Publication Peer Reviews (P³Rs)

One P³R has been posted to this article:
<http://www.pediatrics.org/cgi/eletters/125/2/228>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://www.pediatrics.org/misc/Permissions.shtml>

Reprints

Information about ordering reprints can be found online:
<http://www.pediatrics.org/misc/reprints.shtml>

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

